



## JEE Main Online Exam 2024

### Questions & Solution

04<sup>th</sup> April 2024 | Morning


#### PHYSICS

**Section-A:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct..

- Q.1** An electron is projected with uniform velocity along the axis inside a current carrying long solenoid. Then :
- (1) the electron will be accelerated along the axis.
  - (2) the electron will continue to move with uniform velocity along the axis of the solenoid.
  - (3) the electron path will be circular about the axis.
  - (4) the electron will experience a force at 45° to the axis and execute a helical path.

**Ans.** [2]

**Sol.**



Since  $\vec{v} \parallel \vec{B}$  so force on electron due to magnetic field is zero. So it will move along axis with uniform velocity.

- Q.2** The electric field in an electromagnetic wave is given by  $\vec{E} = \hat{i} 40 \cos \omega \left( t - \frac{z}{c} \right) \text{NC}^{-1}$ . The magnetic field induction of this wave is (in SI unit) :

(1)  $\vec{B} = \hat{i} \frac{40}{c} \cos \omega \left( t - \frac{z}{c} \right)$

(2)  $\vec{B} = \hat{j} 40 \cos \omega \left( t - \frac{z}{c} \right)$

(3)  $\vec{B} = \hat{k} \frac{40}{c} \cos \omega \left( t - \frac{z}{c} \right)$

(4)  $\vec{B} = \hat{j} \frac{40}{c} \cos \omega \left( t - \frac{z}{c} \right)$

**Ans.** [4]

**Sol.**  $\vec{E} = \hat{i} 40 \cos \omega \left( t - \frac{z}{c} \right)$





$\vec{E}$  is along +x direction

$\vec{v}$  is along +z direction

So directional of  $\vec{B}$  will be along +y and magnitude of B will be  $\frac{E}{c}$


So answer is  $\frac{40}{c} \cos \omega \left( t - \frac{z}{c} \right) \hat{j}$

**Q.3** Which of the following nuclear fragments corresponding to nuclear fission between neutron ( ${}^1_0\text{n}$ ) and uranium isotope ( ${}^{235}_{92}\text{U}$ ) is correct :



**Ans.** [4]

**Sol.** Balancing mass number and atomic number



**Q.4** In an experiment to measure focal length ( $f$ ) of convex lens, the least counts of the measuring scales for the position of object ( $u$ ) and for the position of image ( $v$ ) are  $\Delta u$  and  $\Delta v$ , respectively. The error in the measurement of the focal length of the convex lens will be :

(1)  $\frac{\Delta u}{u} + \frac{\Delta v}{v}$

(2)  $f^2 \left[ \frac{\Delta u}{u^2} + \frac{\Delta v}{v^2} \right]$

(3)  $2f \left[ \frac{\Delta u}{u} + \frac{\Delta v}{v} \right]$

(4)  $f \left[ \frac{\Delta u}{u} + \frac{\Delta v}{v} \right]$

**Ans.** [2]

**Sol.**  $f^{-1} = v^{-1} - u^{-1}$

$$-f^{-2} df = -v^{-2} dv - u^{-2} du$$


$$\frac{df}{f^2} = \frac{dv}{v^2} + \frac{du}{u^2}$$

$$df = f^2 \left[ \frac{dv}{v^2} + \frac{du}{u^2} \right]$$

**Q.5** Given below are two statements :

**Statement I :** When speed of liquid is zero everywhere, pressure difference at any two points depends on equation  $P_1 - P_2 = \rho g (h_2 - h_1)$

**Statement II :** In ventury tube shown  $2gh = v_1^2 - v_2^2$




In the light of the above statements, choose the most appropriate answer from the options given below.

- (1) Both Statement I and Statement II are correct.
- (2) Statement I is incorrect but Statement II is correct.
- (3) Both Statement I and Statement II are incorrect.
- (4) Statement I is correct but Statement II is incorrect.

**Ans.** [4]

Sol.



Applying Bernoulli's equation

$$P_1 + \rho gh_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho gh_2 + \frac{1}{2} \rho v_2^2$$

 [ $h_1$  and  $h_2$  are height of point from any reference level]

 Given  $V_1 = V_2 = 0$  (for statement-1)

$$\therefore P_1 - P_2 = \rho g(h_1 - h_2)$$

For statement-2

$$P_1 + \frac{1}{2} \rho v_1^2 = P_2 + \frac{1}{2} \rho v_2^2$$

$$P_1 - P_2 = \rho gh$$

$$P_1 - P_2 = \frac{1}{2} \rho v_2^2 - \frac{1}{2} \rho v_1^2$$

$$\rho gh = \frac{1}{2} \rho v_2^2 - \frac{1}{2} \rho v_1^2$$

$$2gh = v_2^2 - v_1^2$$

Hence answer (4)

**Q.6** The resistances of the platinum wire of a platinum resistance thermometer at the ice point and steam point are  $8 \Omega$  and  $10 \Omega$  respectively. After inserting in a hot bath of temperature  $400^\circ\text{C}$ , the resistance of platinum wire is :

- (1)  $2 \Omega$                                       (2)  $16 \Omega$                                       (3)  $8 \Omega$                                       (4)  $10 \Omega$

Ans. [2]

 Sol. Given  $R_0 = 8 \Omega$ ,  $R_{100} = 10 \Omega$ 

$$\therefore R_{100} = R_0(1 + \alpha \Delta T)$$

$$\text{Also, } R_{400} = R_0(1 + \alpha \Delta T^1)$$

$$\therefore 10 = 8(1 + \alpha \times 100) \Rightarrow 100\alpha = \frac{1}{4}$$

$$\therefore R_{400} = 8(1 + 400\alpha) = 8(1 + 1) = 16 \Omega$$


Hence option (2)

**Q.7** A metal wire of uniform mass density having length  $L$  and mass  $M$  is bent to form a semicircular arc and a particle of mass  $m$  is placed at the centre of the arc. The gravitational force on the particle by the wire is:

- (1)  $\frac{GMm\pi}{2L^2}$                                       (2) 0                                      (3)  $\frac{GmM\pi^2}{L^2}$                                       (4)  $\frac{2GmM\pi}{L^2}$





Ans. [4]

Sol.






**Q.11** A wooden block, initially at rest on the ground, is pushed by a force which increases linearly with time  $t$ . Which of the following curve best describes acceleration of the block with time :



**Ans.** [2]  
**Sol.**



$$F = ma \Rightarrow a = \frac{F}{m} = \frac{kt}{m}$$

$a$  vs  $t$  will be straight line passing through origin.  
Since option (2).

**Q.12** If a rubber ball falls from a height  $h$  and rebounds upto the height of  $h/2$ . The percentage loss of total energy of the initial system as well as velocity ball before it strikes the ground, respectively, are :

- (1) 50%,  $\sqrt{\frac{gh}{2}}$       (2) 50%,  $\sqrt{gh}$       (3) 40%,  $\sqrt{2gh}$       (4) 50%,  $\sqrt{2gh}$

**Ans.** [4]

**Sol.** Velocity just before collision =  $\sqrt{2gh}$

$$\text{Velocity just after collision} = \sqrt{2g\left(\frac{h}{2}\right)}$$

$$\therefore \Delta KE = \frac{1}{2}m(2gh) - \frac{1}{2}mgh = \frac{1}{2}mgh$$

$\therefore$  % loss in energy

$$= \frac{\Delta KE}{KE_1} \times 100 = \frac{\frac{1}{2}mgh}{\frac{1}{2}mg2h} \times 100 = 50\%$$

Hence option (4)

**Q.13** The equation of stationary wave is :  $y = 2a \sin\left(\frac{2\pi nt}{\lambda}\right) \cos\left(\frac{2\pi x}{\lambda}\right)$

Which of the following is NOT correct-

- (1) The dimensions of  $nt$  is [L]      (2) The dimensions of  $n$  is  $[LT^{-1}]$   
(3) The dimensions of  $n/\lambda$  is [T]      (4) The dimensions of  $x$  is [L]

**Ans.** [3]

**Sol.** Comparing the given equation with standard equation of standing  $\frac{2\pi n}{\lambda} = \omega$  and  $\frac{2\pi}{\lambda} = k$

$$\left[ \frac{n}{\lambda} \right] = [\omega] = T^{-1}$$

$$[nt] = [\lambda] = L$$

$$[n] = [\lambda\omega] = LT^{-1}$$

$$[x] = [\lambda] = L$$

Hence option (3)

**Q.14** A body travels 102.5 m in  $n^{\text{th}}$  second and 115.0 m in  $(n + 2)^{\text{th}}$  second. The acceleration is :  
 (1)  $9 \text{ m/s}^2$  (2)  $6.25 \text{ m/s}^2$  (3)  $12.5 \text{ m/s}^2$  (4)  $5 \text{ m/s}^2$

**Ans.** [2]

**Sol.** Given,  $102.5 = u + \frac{a}{2}(2n - 1)$  and  $115 = u + \frac{a}{2}(2n + 3)$

$$\Rightarrow 102.5 = u + an - \frac{a}{2} \text{ and } 115 = u + an + \frac{3a}{2}$$

$$12.5 = 2a \Rightarrow a = 6.25 \text{ m/s}^2$$

Hence option (2)

**Q.15** To measure the internal resistance of a battery, potentiometer is used. For  $R = 10 \Omega$ , the balance point is observed at  $\ell = 500 \text{ cm}$  and for  $R = 1 \Omega$  the balance point is observed at  $\ell = 400 \text{ cm}$ . The internal resistance of the battery is approximately :

- (1)  $0.2 \Omega$  (2)  $0.4 \Omega$  (3)  $0.1 \Omega$  (4)  $0.3 \Omega$

**Ans.** [4]

**Sol.** Let potential gradient be  $\lambda$ .

$$\therefore i \times 10 = \lambda \times 500 = \varepsilon - ir_s$$

$$\Rightarrow 500\lambda = \varepsilon - 50\lambda r_s$$





Also,

$$i' \times 1 = \lambda \times 400 = \varepsilon - i'r_s$$

$$\Rightarrow 400\lambda = \varepsilon - 400\lambda r_s$$


$$\therefore 100\lambda = 350\lambda r_s \Rightarrow r_s = \frac{10}{35} \approx 0.3 \Omega$$

**Q.16** An infinitely long positively charged straight thread has a linear charge density  $\lambda \text{ Cm}^{-1}$ . An electron revolves along a circular path having axis along the length of the wire. The graph that correctly represents the variation of the kinetic energy of electron as a function of radius of circular path from the wire is :



**Ans.** [2]

Sol.



Electric field  $E$  at a distance  $r$  due to infinite long wire is  $E = \frac{2k\lambda}{r}$

For of electron  $\Rightarrow F = eE$

$$F = e \left( \frac{2k\lambda}{r} \right)$$

$$F = \frac{2k\lambda e}{r}$$


This force will provide required centripetal force

$$F = \frac{mv^2}{r} = \frac{2k\lambda e}{r}$$

$$KE = \frac{1}{2}mv^2 = \frac{1}{2}m \left( \frac{2k\lambda e}{m} \right)$$

This is constant so option (2) is correct.

Q.17 The value of net resistance of the network as shown in the given figure is :



(1)  $\left(\frac{5}{2}\right)\Omega$


(2)  $\left(\frac{15}{4}\right)\Omega$

(3)  $6\Omega$

(4)  $\left(\frac{30}{11}\right)\Omega$

Ans. [3]

Sol.




Diode 2 is in reverse bias

So current will not flow in branch of 2<sup>nd</sup> diode, So we can assume it to be broken wire.

Diode 1 is in forward bias


So it will behave like conducting wire. So new circuit will be



$$R_{eq} = \frac{15 \times 10}{15 + 10} = \frac{15 \times 10}{25} = 6 \Omega$$

Correct answer (3)

**Q.18** P-T diagram of an ideal gas having three different densities  $\rho_1, \rho_2, \rho_3$  (in three different cases) is shown in the figure. Which of the following is correct :



(1)  $\rho_2 < \rho_3$

(2)  $\rho_1 > \rho_2$

(3)  $\rho_1 < \rho_2$

(4)  $\rho_1 = \rho_2 = \rho_3$

**Ans.**

[2]

**Sol.**

For ideal gas


$$PV = nRT$$

$$PV = \frac{m}{M} RT$$

$$P = \left( \frac{m}{V} \right) \frac{RT}{M}$$

$$P = \frac{\rho RT}{M}$$

(Where  $m$  is mass of gas and  $M$  is molecular mass of gas)



for same temperature  $P_1 > P_2 > P_3$

So  $\rho_1 > \rho_2 > \rho_3$

So correct answer is (2)

**Q.19** The co-ordinates of a particle moving in x-y plane are given by :

$$x = 2 + 4t, y = 3t + 8t^2.$$

The motion of the particle is :

- (1) non-uniformly accelerated.
- (2) uniformly accelerated having motion along a straight line.
- (3) uniform motion along a straight line.
- (4) uniformly accelerated having motion along a parabolic path.

**Ans.**

[4]

**Sol.**

$$x = 2 + 4t$$

$$\frac{dy}{dt} = v_x = 4$$



$$\frac{dv_x}{dt} = a_x = 0$$

$$y = 3t + 8t^2$$

$$\frac{dy}{dt} = v_y = 3 + 16t$$

$$\frac{dv_y}{dt} = a_y = 16$$

the motion will be uniformly accelerated motion.

For path

$$x = 2 + 4t$$

$$\frac{(x-2)}{4} = t$$

Put this value of t is equation of y

$$y = 3\left(\frac{x-2}{4}\right) + 8\left(\frac{x-2}{4}\right)^2$$

this is a quadratic equation so path will be parabola.

Correct answer (4).

**Q.20** In an ac circuit, the instantaneous current is zero, when the instantaneous voltage is maximum. In this case, the source may be connected to :

- A. pure inductor.
- B. pure capacitor.
- C. pure resistor.
- D. combination of an inductor and capacitor.

Choose the correct answer from the options given below :

- (1) A, B and C only
- (2) B, C and D only
- (3) A and B only
- (4) A, B and D only

**Ans.** [4]


**Sol.** This is possible when phase difference is  $\frac{\pi}{2}$  between current and voltage so correct answer will be (4)

**Section-B: Numerical Value Type Questions:** This section contains 10 Numerical based questions. Attempt any 5 questions out of 10. The answer to each question should be rounded-off to the nearest integer.

**Q.21** An infinite plane sheet of charge having uniform surface charge density  $+\sigma_s$  C/m<sup>2</sup> is placed on x-y plane. Another infinitely long line charge having uniform linear charge density  $+\lambda_c$  C/m is placed at  $z = 4$  m plane and parallel to y-axis. If the magnitude value  $|\sigma_s| = 2|\lambda_c|$  then at point  $(0, 0, 2)$ , the ratio of magnitudes of electric field values due to sheet charge to that of line charge is  $\pi\sqrt{n} : 1$ . The value of n is \_\_\_\_\_.

**Ans.** [16]

**Sol.**



$$\begin{aligned} \frac{E_s}{E_\ell} &= \frac{\sigma}{2\epsilon_0} \times \frac{2\pi\epsilon_0 r}{\lambda} \\ &= \frac{\pi \times \sigma r}{\lambda} \\ &= \frac{\pi \times 2\lambda \times 2}{\lambda} = \frac{4\pi}{1} \end{aligned}$$

$$\therefore n = 16$$

**Q.22** A hydrogen atom changes its state from  $n = 3$  to  $n = 2$ . Due to recoil, the percentage change in the wave length of emitted light is approximately  $1 \times 10^{-n}$ . The value of  $n$  is \_\_\_\_\_. [Given  $Rhc = 13.6$  eV,  $hc = 1242$  eV nm,  $h = 6.6 \times 10^{-34}$  J s, mass of the hydrogen atom =  $1.6 \times 10^{-27}$  kg]

**Ans.** [7]

**Sol.**  $\Delta E = 13.6 \left( \frac{1}{2^2} - \frac{1}{3^2} \right) = 1.9$  eV

$$\Delta E = \frac{hc}{\lambda}$$

$$\lambda = \frac{hc}{\Delta E}$$

$$P_i = P_f$$

$$0 = -mv + \frac{h}{\lambda'} \Rightarrow v = \frac{h}{m\lambda'}$$

$$\Delta E = \frac{1}{2}mv^2 + \frac{hc}{\lambda'}$$

$$= \frac{1}{2}m \left( \frac{h}{m\lambda'} \right)^2 + \frac{hc}{\lambda'}$$

Now

$$\Delta E = \frac{h^2}{2m\lambda'^2} + \frac{hc}{\lambda'}$$

$$\lambda'^2 \Delta E - hc\lambda' - \frac{h^2}{2m} = 0$$

$$\lambda' = \frac{hc \pm \sqrt{h^2c^2 + \frac{4\Delta E h^2}{2m}}}{2\Delta E}$$

$$\lambda' = \frac{hc \pm hc \sqrt{1 + \frac{2\Delta E}{mc^2}}}{2\Delta E}$$

$$\frac{\lambda'}{\lambda} = \frac{1 + \left(1 + \frac{2\Delta E}{mc^2}\right)^{1/2}}{2} = \frac{1 + 1 + \frac{\Delta E}{mc^2}}{2}$$


$$\frac{\lambda'}{\lambda} = 1 + \frac{\Delta E}{mc^2}$$

$$\frac{\lambda' - \lambda}{\lambda} = \frac{\Delta E}{2mc^2} = \frac{1.9 \times 1.6 \times 10^{-19}}{2 \times 1.67 \times 10^{-27} \times 9 \times 10^{16}} = 10^{-9}$$

$$\therefore \% \text{ change} \approx 10^{-7}$$

Correct answer 7.

- Q.23** The magnetic field existing in a region is given by  $\vec{B} = 0.2(1 + 2x)\hat{k}T$ . A square loop of edge 50 cm carrying 0.5A current is placed in x-y plane with its edge parallel to the x-y axes, as shown in figure. The magnitude of the net magnetic force experienced by the loop is \_\_\_\_\_ mN.



**Ans.** [50]

**Sol.** Force on segment parallel to x-axis will cancel each other. Hence  $F_{\text{net}}$  will be due to portion parallel to y-axis.

$$\begin{aligned} F &= 0.5 \times 0.5 \times 6 \times 0.2 - 0.5 \times 0.5 \times 0.2 \times 5 \\ &= 0.5 \times 0.5 \times 0.2 \\ &= 0.25 \times 0.2 \\ &= 50 \times 10^{-3} \text{ N} \\ &= 50 \text{ mN} \end{aligned}$$


- Q.24** A alternating current at any instant is given by  $i = \left[ 6 + \sqrt{56} \sin\left(100\pi t + \frac{\pi}{3}\right) \right]$  A. The rms value of the current is \_\_\_\_\_ A.

**Ans.** [8]

**Sol.** 
$$I_{\text{rms}} = \sqrt{\frac{\int i^2 dt}{\int dt}}$$


$$I_{\text{rms}} = \sqrt{(6)^2 + \frac{(\sqrt{56})^2}{2}} = \sqrt{36 + 28} = \sqrt{64} = 8 \text{ A}$$

- Q.25** Twelve wires each having resistance  $2\Omega$  are joined to form a cube. A battery of 6 V emf is joined across point a and c. The voltage difference between e and f is \_\_\_\_\_ V.



**Ans.** [1]

**Sol.**



For symmetry, current through e-b and g-d = 0

$$\therefore R_{\text{eq}} = \frac{4}{3} \times R = \frac{3}{2} \Omega$$

$$\therefore \text{Current through battery} = \frac{6 \times 2}{3} = 4 \text{ A}$$

$$i_2 = \frac{4}{8} \times 2 = 1 \text{ A}$$

$$\therefore \Delta V \text{ across e-f} = \frac{i_2}{2} \times R = \frac{1}{2} \times 2 = 1 \text{ V}$$

**Q.26** A soap bubble is blown to a diameter of 7 cm. 36960 erg of work is done in blowing it further. If surface tension of soap solution is 40 dyne/cm then the new radius is \_\_\_\_\_ cm. Take  $\left(\pi = \frac{22}{7}\right)$

**Ans.** [7]

**Sol.**  $W = \Delta U = S\Delta A$

$$36960 \text{ erg} = \frac{40 \text{ dyne}}{\text{cm}} 8\pi \left[ (r)^2 - \left(\frac{7}{2}\right)^2 \right] \text{ cm}^2$$

$$r = 7 \text{ cm}$$

**Q.27** Two wavelengths  $\lambda_1$  and  $\lambda_2$  are used in Young's double slit experiment  $\lambda_1 = 450 \text{ nm}$  and  $\lambda_2 = 650 \text{ nm}$ . The minimum order of fringe produced by  $\lambda_2$  which overlaps with the fringe produced by  $\lambda_1$  is n. The value of n is \_\_\_\_\_.

**Ans.** [9]

**Sol.**  $n_2 \lambda_2 = n_1 \lambda_1$

$$\frac{n_2}{n_1} = \frac{\lambda_1}{\lambda_2}$$

$$= \frac{450}{650} = \frac{9}{13}$$

$$n_2 = 9$$

**Q.28** An elastic spring under tension of 3 N has a length a. Its length is b under tension 2 N. For its length  $(3a - 2b)$ , the value of tension will be \_\_\_\_\_ N.

**Ans.** [5]

**Sol.**  $3 = K(a - \ell)$

$$2 = K(b - \ell)$$

$$T = K(3a - 2b - \ell)$$

$$T = K(3(a - \ell) - 2(b - \ell))$$

$$= K \left[ 3 \left( \frac{3}{K} \right) - 2 \left( \frac{2}{K} \right) \right]$$

$$= 9 - 4$$

$$= 5 \text{ N}$$

**Q.29** Two forces  $\vec{F}_1$  and  $\vec{F}_2$  are acting on a body. One force has magnitude thrice that of the other force and the resultant of the two forces is equal to the force of larger magnitude. The angle between  $\vec{F}_1$  and  $\vec{F}_2$  is  $\cos^{-1}\left(\frac{1}{n}\right)$ . The value of  $|n|$  is \_\_\_\_\_.

**Ans.** [6]

**Sol.**  $|\vec{F}_1| = F$

$$|\vec{F}_R| = |\vec{F}_2| = 3F$$

$$\vec{F}_R^2 = F_1^2 + F_2^2 + 2F_1F_2 \cos \theta$$

$$9F^2 = F^2 + 9F^2 + 6F^2 \cos \theta$$

$$\cos \theta = -\frac{1}{6}$$

$$\theta = \cos^{-1}\left(-\frac{1}{6}\right)$$

$$n = -6$$

$$|n| = 6$$

**Q.30** A solid sphere and a hollow cylinder roll up without slipping on same inclined plane with same initial speed  $v$ . The sphere and the cylinder reaches upto maximum heights  $h_1$  and  $h_2$ , respectively, above the initial level.

The ratio  $h_1 : h_2$  is  $\frac{n}{10}$ . The value of  $n$  is \_\_\_\_\_.

**Ans.** [7]

**Sol.** Gain in P.E. = Loss in K.E.

$$mgh = \frac{1}{2}mv^2\left(1 + \frac{K^2}{R^2}\right)$$

$$h \propto 1 + \frac{K^2}{R^2}$$

$$\frac{h_1}{h_2} = \frac{1 + \frac{2}{5}}{1 + 1} = \frac{7}{5 \times 2} = \frac{7}{10}$$

$$n = 7$$

**CHEMISTRY**

**Section-A:** This section contains 20 multiple choice questions. Each question has 4 choices(1), (2), (3) and (4), out of which **ONLY ONE** is correct..

**Q.31** What pressure (bar) of  $H_2$  would be required to make emf of hydrogen electrode zero in pure water at  $25^\circ C$  ?  
 (1)  $10^{-14}$  (2)  $10^{-7}$  (3) 1 (4) 0.5

**Ans.** [1]

**Sol.**  $2e^- + 2H^+(aq) \rightarrow H_2(aq)$

$$E = E^\circ - \frac{0.059}{n} \log \frac{P_{H_2}}{[H^+]^2}$$

$$0 = 0 - \frac{0.059}{2} \log \frac{P_{H_2}}{(10^{-7})^2}$$

$$\log \frac{P_{H_2}}{(10^{-7})^2} = 0$$

$$\frac{P_{H_2}}{10^{-14}} = 1$$

**Q.32** The correct sequence of ligands in the order of decreasing field strength is :

(1)  $CO > H_2O > F^- > S^{2-}$

(2)  $^-OH > F^- > NH_3 > CN^-$

(3)  $NCS^- > EDTA^{4-} > CN^- > CO$

(4)  $S^{2-} > ^-OH > EDTA^{4-} > CO$

**Ans.** [1]

**Sol.** According to spectrochemical series ligand field strength is  $CO > H_2O > F^- > S^{2-}$

**Q.33** Match List -I with List II:


	<b>List - I</b> <b>Mechanism steps</b>	<b>List - II</b> <b>Effect</b>
(A)		(I) - E effect
(B)		(II) - R effect
(C)		(III) + E effect
(D)		(IV) + R effect

Choose the **correct** answer from the options given below :



Due to common ion effect of  $\text{NH}_4^+$ ,


$[\text{OH}^-]$  decreases in such extent that only group-III cation can be precipitated, due to their very low  $K_{sp}$  in the range of  $10^{-38}$ .



Identify (B) and (C) and how are (A) and (C) related ?

	(B)	(C)	
(1)			functional group isomers
(2)			Derivative
(3)			position isomers
(4)			chain isomers

Ans. [3]  
Sol.



A and C are position isomer.

















































